## Mathematical analysis II Homework 1

To be handed in by Wednesday, 15.10.25, 23:59 h via OWL

## Exercise 1 (Metrics).

(3+1=4)

a) Let  $X = [1, \infty)$ . Show that the function defined via

$$d(x,y) = \left| \frac{1}{x} - \frac{1}{y} \right|$$

is a metric.

b) Let  $X = \mathbb{R}$ . Is the function defined by

$$d(x,y) = |x|^{|y|}$$

a metric? Show or disprove!

**Solution.** a) Symmetry and  $d(x,y) \ge 0$  is obvious since the absolute value  $|\cdot|$  fulfils this. If d(x,y) = 0, then

$$\left|\frac{1}{x} - \frac{1}{y}\right| = 0 \Leftrightarrow \frac{1}{x} - \frac{1}{y} = 0 \Leftrightarrow \frac{1}{x} = \frac{1}{y} \Leftrightarrow x = y,$$

where the first equivalence uses that |a| = 0 if and only if a = 0. Finally, the triangle inequality follows by

$$d(x,y) = \left| \frac{1}{x} - \frac{1}{y} \right| = \left| \frac{1}{x} - \frac{1}{z} + \frac{1}{z} - \frac{1}{y} \right| \le \left| \frac{1}{x} - \frac{1}{z} \right| + \left| \frac{1}{z} - \frac{1}{y} \right| = d(x,z) + d(z,y),$$

and the inequality comes from the triangle inequality fulfilled by  $|\cdot|$ . (Note that inserting  $\pm 1/z$  corresponds to adding a zero, something that every mathematician should be able to do.) Note also that even if X for us is an infinite axis, the so-called diameter of this space X under d is

$$\operatorname{diam}(X) := \sup\{d(x,y) : x,y \in X\} = 1$$

such that X is bounded and equal to  $B_1^d$ , its ball of radius 1.

*Remark:* Many of you did this wrongly, so let me emphasize that definiteness means  $d(x,y) = 0 \Rightarrow x = y$ . That d(x,x) = 0 is usually trivial to see.

b) It is not a metric since it violates (almost) everything: first, it is not symmetric by setting, e.g., x = 1 and y = 2. Second, it is not definite since for any  $y \neq 0$ , we have  $d(0, y) = |0|^{|y|} = 0$ . Third, triangle inequality fails, set x = 3, y = 2, z = 1.

## Exercise 2 (Open and closed sets).

(3+3=6)

In here, we assume  $X = \mathbb{R}$ .

a) Let  $d: X \times X \to [0, \infty)$  be a metric, and define

$$\delta(x, y) = \min\{d(x, y), 1\}.$$

Show that  $\delta$  is really a metric, and that d and  $\delta$  define the same open and closed sets.

b) Let

$$d(x,y) = \begin{cases} 1 & \text{if } x \neq y, \\ 0 & \text{if } x = y. \end{cases}$$

Show that under this metric, every set is both open and closed.

**Solution.** a) Obviously  $\delta \geq 0$  since both  $d(x,y) \geq 0$  and  $1 \geq 0$ . Moreover

$$\delta(x,y) = 0 \Leftrightarrow \min\{d(x,y),1\} = 0 \Leftrightarrow d(x,y) = 0 \Leftrightarrow x = y$$

since d is a metric. Symmetry is trivial as well since d(x, y) is symmetric as it is a metric. Triangle inequality follows from

$$\delta(x,y) = \min\{d(x,y), 1\} \le \min\{d(x,z) + d(z,y), 1\} \le \min\{d(x,z), 1\} + \min\{d(z,y), 1\},$$

where the first inequality uses that d is a metric and that  $\min\{a, c\} \leq \min\{b, c\}$  for any  $a \leq b$  and any c, and the last equality uses subadditivity of min for non-negative arguments (for understanding: find an example where  $\min\{a+b,1\} > \min\{a,1\} + \min\{b,1\}$ ).

Let's come to open sets. I will denote

$$B_{\varepsilon}^d(a) = \{x \in X : d(x, a) < \varepsilon\}, \qquad B_{\varepsilon}^{\delta}(a) = \{x \in X : \delta(x, a) < \varepsilon\}.$$

We have to show 2 directions:

1) Let  $A \subset X$  be open wrt.  $\delta$ , that is,

$$\forall a \in A \exists \varepsilon > 0 : B_{\varepsilon}^{\delta}(a) \subset A.$$

Now, we need to find for any  $a \in A$  some  $\varepsilon_1 > 0$  such that  $B^d_{\varepsilon_1}(a) \subset A$ . This is achieved by  $\varepsilon_1 = \varepsilon$ , since by definition, always  $\delta(x,y) \leq d(x,y)$  for any  $x,y \in X$ , hence  $B^d_{\varepsilon}(a) \subset B^\delta_{\varepsilon}(a)$  for any a (notice the reversed subset relation). In other words, if  $B^\delta_{\varepsilon}(a) \subset A$ , then also  $B^d_{\varepsilon}(a) \subset B^\delta_{\varepsilon}(a) \subset A$ , and hence A is open wrt. d.

2) Let  $A \subset X$  be open wrt. d, that is,

$$\forall a \in A \exists \varepsilon > 0 : B_{\varepsilon}^{d}(a) \subset A.$$

Again, we need to find for any  $a \in A$  some  $\varepsilon_1 > 0$  such that  $B^{\delta}_{\varepsilon_1}(a) \subset A$ . This is achieved by  $\varepsilon_1 = \min\{\varepsilon, 1\}$ . Indeed, if  $x \in B^{\delta}_{\varepsilon_1}(a)$ , then  $\delta(x, a) < \varepsilon_1 = \min\{\varepsilon, 1\} \leq 1$ , which means that  $\delta(x, a) = d(x, a)$  (otherwise we would have  $\delta(x, a) = 1$ , which contradicts the fact that  $\delta(x, a) < 1$ ). But then  $d(x, a) = \delta(x, a) < \min\{\varepsilon, 1\} \leq \varepsilon$ , i.e.,  $d(x, a) < \varepsilon$ , giving  $x \in B^d_{\varepsilon}(a)$  and so  $x \in A$  as A is open wrt. d. This shows that for this  $\varepsilon_1$ , we have  $B^{\delta}_{\varepsilon_1}(a) \subset A$  and thus A is open wrt.  $\delta$ .

Finally, the also define the same closed sets: let  $A \subset X$  be closed wrt. d, then by definition,  $X \setminus A$  is open under d. But now we just showed that this means that  $X \setminus A$  is open under  $\delta$ , and again according to the definition, A closed wrt.  $\delta$ . Obviously this holds also if we interchange d and  $\delta$ . This ends the proof.

b) Let  $A \subset X$ . First we show that A is open, that means for any  $a \in A$ , we need to find  $\varepsilon > 0$  such that  $B_{\varepsilon}(a) \subset A$ . We choose (amazingly independently of the set A!)  $\varepsilon = \frac{1}{2}$ , then, by definition of d, we have  $B_{\varepsilon}(a) = \{a\}$ , a single point. Since  $a \in A$ , this means  $B_{\varepsilon}(a) = \{a\} \subset A$  such that A is open. Next, A is also closed: by definition (or one of the equivalent definitions), A is closed if and only if  $X \setminus A$  is open. Since obviously  $X \setminus A \subset X$  is a subset, and we just showed that any subset of X is open, also  $X \setminus A$  is open. According to the definition, this means that A is closed. (Here, we could also argue with sequences, since under d any convergent sequence will be constant from some point on. Try to formalize this!)